

Henry Ford Health System 6777 W. Maple Rd., West Bloomfield, MI 48322 2825 Livernois, Troy, MI 48083 Ph: 248-952-9205

www.ericmakhnimd.com

PHYSICAL THERAPY PROTOCOL: Distal Femoral Osteotomy

Name:	Date of Surgery:
Diagnosis:	Signed:

Frequency: 2-3 times per week x 12 weeks

Phase $1 \rightarrow 0-2$ Weeks Post-Op

Weight Bearing:

- Heel touch weight bearing with brace on

Brace:

On at all times, day and night, off for shower and exercises only

Range of Motion:

- 0 to 90 degrees at home

Exercises:

- Calf pumps, quad sets, SLR in brace, modalities per PT discretion

Phase $2 \rightarrow 2-6$ Weeks Post-Op

- Heel touch weight bearing with brace
- Open from 0 to 90 degrees and wear until 6 weeks
- Progress flexion as tolerated until full ROM

Exercises:

- Progress non-weight bearing flexibility, modalities per PT discretion
- Begin floor based core and glute exercises, advance quad sets, patellar mobility, and SLR

Phase $3 \rightarrow 6-8$ Weeks Post-Op

-d/c brace

Weight Bearing:

- Advance 25% each week until full weight bearing and normalized gait pattern

Exercises:

- Advance closed chain quads, progress balance, core/pelvic and stability work
- Start stationary bike at 6 weeks, advance SLR and floor based exercises

Phase $4 \rightarrow 8$ Weeks-4 Months Post-Op

- Full weight bearing/as tolerated

Exercises:

- Progress flexibility/strengthening, progression of functional balance, core, and glutes program
- Advance bike, add elliptical at 12 weeks as tolerated, may swim at 12 weeks

Phase $5 \rightarrow 4-6$ Months Post-Op

Exercises:

- Advance Phase 4 activities
- Progress to functional training, including impact activity after 20 wks when cleared by MD