

Henry Ford Health System 6777 W. Maple Rd., West Bloomfield, MI 48322 2825 Livernois, Troy, MI 48083 Ph: 248-661-7029

www.ericmakhnimd.com

PHYSICAL THERAPY PROTOCOL/PRESCRIPTION: TOTAL SHOULDER REPLACEMENT

Name:	Date of Surgery:
Diagnosis:	Signed:
Frequency: 2-3 times per week x 6 weeks.	
Weeks 1-6:	

- PROM AAROM as tolerated, except the following. . .
 - No active IR/backwards extension for 6 weeks. The subscapularis tendon is taken down for the surgery and then repaired afterwards. It takes about 4-6 weeks for it to grown back into the humerus and regenerate a blood and nerve supply.
- PROM in biceps flexion for first 6 weeks
- ROM goals: Week 1: 120° FF/20° ER at side; ABD max 75° without rotation
- ROM goals: Week 2: 140° FF/40° ER at side; ABD max 75° without rotation
- No resisted internal rotation/backward extension until 12 weeks post-op
- Grip strengthening OK
- Canes/pulleys OK if advancing from PROM
- Heat before PT, ice after PT

Weeks 6-12:

- Begin AAROM AROM for internal rotation and backwards extension as tolerated, if not already begun.
- Goals: Increase ROM as tolerated with gentle passive stretching at end ranges
- Begin light resisted ER/FF/ABD: isometrics and bands, concentric motions only
- No resisted internal rotation/backwards extension until 12 weeks post-op
- No scapular retractions with bands yet

Months 3-12:

- Begin resisted IR/BE (isometrics/bands): isometrics à light bands à weights
- Advance strengthening as tolerated; 10 reps/1 set per exercise for rotator cuff, deltoid, and scapular stabilizers.
- Increase ROM to full with passive stretching at end ranges
- Begin eccentric motions, plyometrics, and closed chain exercises at 12 weeks.