

Henry Ford Health System 6777 W. Maple Rd., West Bloomfield, MI 48322 2825 Livernois, Troy, MI 48083

Ph: 248-661-7029 www.ericmakhnimd.com

# PHYSICAL THERAPY PROTOCOL: Hip Gluteus Medius Repair (w/ or w/out labral debridement)

Name:	Date of Surgery:
	- '
Diagnosis:	Signed:

# Frequency: 2-3 times per week x 12 weeks

# Phase $1 \rightarrow 0-4$ Weeks Post-Op

- Partial weight bearing for 6 weeks with walker
- CPM for 4 hours/day

## Range of motion:

- PROM
  - o Hip flexion to 90 degrees, abduction as tolerated
  - o No active abduction and IR
  - No passive ER or adduction for 6 weeks

#### Exercises:

- Bike for 20 minutes/day, can be 2x/day
- Quaddruped rocking for hip flexion
- Hip isometrics → extension, adduction, ER at 2 weeks
- Pelvic tilts
- NMES to quads with SAQ

Modalities per PT discretion

## Phase $2 \rightarrow 4-6$ Weeks Post-Op

- Weight bearing 20 lbs through 6 weeks, gait training PWB with walker
- Progress passive hip flexion past 90 degrees

## **Exercises:**

- Continue previous phase exercises
- Supine bridges
- Isotonic adduction
- Progress core strengthening (avoid hip flexor tendonitis)
- Progress hip strengthening
  - Start isometric sub max pain free hip flexion (3-4weeks)
  - Quadriceps strengthening
- Agua therapy in low end of pool

Modalities per PT discretion

# Phase $3 \rightarrow 6-8$ Weeks Post-Op

- Increase weight bearing to full by 8 weeks with crutches

#### Exercises:

- Continue previous phase exercises
- Progress ROM, passive hip ER/IR
- Supine log rolling → Stool rotation → Standing on BAPS
  - Hip joint mobs with mobilization belt if needed
- Lateral and inferior rotation
- Prone posterior-anterior glides with rotation
  - o Progress core strengthening (avoid hip flexor tendonitis)

# Phase 4 → 8-10 Weeks Post-Op

Wean off crutches

## Exercises:

- Continue previous phase exercises
- Progress hip ROM
- Progress strengthening LE
  - Hip isometrics for abduction and progress to isotonics
  - Leg press (bilateral LE)
  - o Isokinetics: knee flexion/extension
- Progress core strengthening
- Begin proprioception/balance
  - o Balance board and single leg stance
- Bilateral cable column rotations
- Elliptical

# Phase $5 \rightarrow 10-12$ Weeks Post-Op

#### Exercises:

- Continue previous phase exercises
- Progress hip ROM
- Progress strengthening LE and core
  - o Hip PREs and hip machine
  - o Unilateral leg press, Unilateral cable column rotations
  - Hip hiking, Step downs
- Hip flexor, glute/piriformis, and IT band stretching---manual and self
- Progress balance and proprioception
- Bilateral  $\rightarrow$  unilateral  $\rightarrow$  foam  $\rightarrow$  dynadisc
- Treadmill side stepping from level surface holding on progressing to inclines
- Side stepping with theraband
- Hip hiking on stairmaster (week12)

## Phase $5 \rightarrow 12$ Weeks Post-Op

## **Exercises:**

- Progress hip ROM and stretching, progress LE and core strengthening
- Endurance activities around the hip
- Dynamic balance activities
- Treadmill running program
- Sport specific agility drills and plyometrics
- D/C criteria (3-6 months)
  - Hip outcome score, pain free or manageable level of discomfort, MMT within 10% of non-op limb, biodex test of quad and hamstrings peak torque within 15% of non-op limb, single leg cross-over triple hop for distance: (less than 85% is abnormal), step down test